

Free Fall / review problems

Dude drops water balloon at $t=0$

from height $y_0 = (?)$. Given: $a = 9.8$ down

$y_1 = 50, y_2 = 30$
 $v_1 = -20$ 3.14, E

(a) Find v_2


Soln: $v_2^2 - v_1^2 = 2a(y_2 - y_1) \leftarrow (2.11)$

with $v_1 = -20, a = -9.8, y_2 - y_1 = 30 - 50 = -20$

$$v_2^2 - 20^2 = 2(-9.8)(-20)$$

$$v_2^2 = 400 + 392$$

$$v_2 = \sqrt{792} = \boxed{28.14 \text{ m/s}}$$



b) Assuming $v_0 = 0$, what was y_0 ?

Soln: $v_1^2 - v_0^2 = 2a_y(y_1 - y_0) \quad (E)$

$$(-20)^2 - 0 = 2(-9.8)(50 - y_0)$$

$$400 = -19.6 \times 50 + 19.6 y_0$$

$$\frac{400 + 980}{19.6} = y_0$$

$$\boxed{y_0 = 70.4 \text{ m}}$$

c) How long does it take to splash down?

$$y(t_{\text{splash}}) = y_0 + v_{0y}t_s + \frac{1}{2}at_s^2 \quad (3.13)$$

$$0 = 70.4 + 0 - \frac{9.8}{2}t_s^2$$

$$-70.4 = -4.9t_s^2$$

$$14.4s = t_s^2 \Rightarrow \boxed{t_s = 3.79 \text{ sec}}$$

Note: when $v_{0y} \neq 0$, use quadratic formulas

$$at^2 + bt + c = 0$$

$$(y - y_0)$$

$$t = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

d) How fast is it moving at t_s ?

$$v_y(t_s) = v_{0y} + a_y t_s \quad \leftarrow (2.13)$$

$$v_y(t_s) = 0 + -9.8(3.79)$$

$$\boxed{v(t_s) = 37.15 \text{ m/s}}$$