## Physics 231. Mechanics of Solids

Exam-like questions - Ch. 4-5. Newton's laws and forces

- 1. Three strings exert forces on a frictionless, dry ice block simulataneously. String 1 pulls the block towards  $\theta = 300^{\circ}$ . String 2 pulls it towards  $\theta = 210^{\circ}$ . String 3 pulls it towards  $\theta = 120^{\circ}$ . Each string pulls with a force of 2 N. What is the net magnitude and direction of the force on the block? (Hint: draw the picture first!)
  - (a) 2N at 300° (b) 1.8N at 190° (c) 2N at 210° (d) 1.8N at 230° (e) 2N at 120°
- 2. A space ranger weighs himself on the Moon, where  $g_{Moon} = 1.625 \text{ m/s}^2$ ?, and the scale reads 243.75 N. What is the mass of the ranger on Earth (wearing the same gear)? Mass = \_\_\_\_\_,
  - (a) 148 N (b) 150 kg (c) 1470 N (d) 400 kg (e) 25 kg
- 3. A space ranger weighs himself on the Moon, where  $g_{Moon} = 1.625 \text{ m/s}^2$ ?, and the scale reads 243.75 N. What is the weight of the ranger on Earth (wearing the same gear)? Weight = \_\_\_\_\_.
  - (a) 1470 N (b) 148 kg (c) 400 N (d) 90 kg (e) 880 N
- 4. A 90 kg man travels in an elevator accelerating downward at 2 m/ $s^2$ . What force is exerted on the man by the elevator floor?
  - (a) 1060 N (b) 880 N (c) 800 N (d) 750 N (e) 700 N
- 5. An elevator with a mass of 2042 kg is supported by a steel cable. What is the tension in the cable when the elevator is being accelerated upward at a rate of 2.0 m/s<sup>2</sup>? (g=9.8 m/s<sup>2</sup>)
  - (a) 10,005 N (b) 23,100 N (c) 24,100 N (d) 26,000 N (e) 40,020 N
- 6. What is the *total acceleration* vector for a particle moving at 5 m/s in a circle of radius 3 m while speeding up at  $3 \text{ m/s}^2$ ?
  - (a)  $-8.33\hat{r} + 3\hat{\theta}$  (b)  $-1.67\hat{r} + 3\hat{\theta}$  (c)  $-6.67\hat{r} 1\hat{\theta}$  (d)  $-8.33\hat{r} + 1\hat{\theta}$ (e)  $1.67\hat{r} + 3\hat{\theta}$
- 7. The force that resists the relative motion of two objects that are sliding against each other is called \_\_\_\_\_\_.
  - (a) static friction
    (b) dynamic friction
    (c) kinetic friction
    (d) rolling friction
    (e) tangential force

- 8. A 30 kg box is pulled horizontally with a uniform force of 181 N. If it starts at rest, what will be its acceleration? (The materials have friction with  $\mu_s = 0.6$  and  $\mu_k = 0.4$ .)
  - (a)  $0 \text{ m/s}^2$  (b)  $0.2 \text{ m/s}^2$  (c)  $1.3 \text{ m/s}^2$  (d)  $2.1 \text{ m/s}^2$  (e)  $5.9 \text{ m/s}^2$
- 9. A 30 kg box is pulled horizontally with a uniform force of 170 N. If it starts at rest, what will be its acceleration? (The materials have friction with  $\mu_s = 0.6$  and  $\mu_k = 0.4$ .)
  - (a)  $0 \text{ m/s}^2$  (b)  $0.2 \text{ m/s}^2$  (c)  $1.3 \text{ m/s}^2$  (d)  $2.1 \text{ m/s}^2$  (e)  $5.9 \text{ m/s}^2$
- 10. A block slides down a ramp at constant speed. The ramp is inclined at  $\theta$  degrees relative to the horizontal. The  $\mu_k$  for the ramp-block system must be \_\_\_\_\_.
  - (a) g/m (b) gsin  $\theta$  (c) 1.0 (d)  $\frac{\cos \theta}{\sin \theta}$  (e) tan  $\theta$
- 11. An automobile moves on a level horizontal road in a circle of radius 30 m. The coefficient of friction between tires and road is 0.50. The maximum speed with which this car can round this curve (without slipping) is:
  - (a) 3.0 m/s (b) 4.9 m/s (c) 9.8 m/s (d) 12. m/s (e) 13. m/s
- 12. A pendulum bob of mass m is offset by the angle  $\theta$  from the vertical at the moment its speed is v = along on arc of radius 2 m. The tension in the string is given by \_\_\_\_\_
  - (a) mg (b)  $mg\sin\theta + mv^2/r$  (c)  $mg\cos\theta + mv^2/r$  (d)  $mv^2/r$ (e)  $mg\tan\theta$