PHYS2321
Week4: (Finish Gauss' Law) and Electric Potential

Day 2 Outline

1) Hwk: Ch. 23 P. 2,3,5,9,12,15,17,21,25,28,29,35,36,43,
48,51. MCQ 1-13 odd (Due Mon)
Read Ch. 23-1 to 23-8
2) Use Gauss' law for nested conducting spheres
3) Quiz 2 on Gauss' law
4) Electric Potential
Electric Potential Energy U_E
Comparison to gravity
Electric potential (or voltage) V
Relation to E-field



Notes: Return Ch. 22 Hwk 3 Mean=7.9/10. Checked #10,35, MQ.
PDF version of this week4 PPT online.
Tutoring Thu 7 pm.

Electric Potential of a Point Charge

- Recall E field

$$\vec{E} = \frac{kq}{r^2}\hat{r}$$

- Electric Potential of a point charge

$$V = \frac{kq}{r}$$

$V = 0 \quad \text{when} \quad r \rightarrow \infty$



- Electric potential difference

$$\Delta V_{ab} = \frac{kq}{r_b} - \frac{kq}{r_a}$$

PHYS2321
Week4: Electric Potential

Day 3 Outline

1) Hwk: Ch. 23 P. 2,3,5,9,12,15,17,21,25,28,29,35,36,43, 48,51. MCQ 1-13 odd (Due Mon)
   Read Ch. 23-1 to 23-8
2) Return Quiz 2. = 3.9/8
3) Electric Potential and Electric Potential energy
   - Comparison to gravity
   - Calculating V for point charges
   - Calculating U for point charges
   - Equipotentials



Notes:

PDF version of this week4 PPT online.

Try “Ch. 23 Test Bank Practice” online.

Electric Potential

- Electric Potential of a point charge (last slide)
- Electric potential closely related to *potential energy*
  - $\Delta U = q\Delta V$
  - And to *work*: $W_{byfield} = -q\Delta V = -\Delta U$
  - Convention: both U and V = 0 at r=infinity
- Electric potential closely related to electric force
  - $F_E \Delta r = W_{byfield} = -q\Delta V$
- Electric potential closely related to electric field
  - $\delta V = -E\delta r$ so that potential difference is: $\Delta V = -\int \vec{E} \cdot d\vec{l}$
- Electric potential is easier to find than the E-field because it is not a vector



Analogy with gravity

F_E on q_0 is down

$W_{Efield} = -|F_E|d \quad (F_E = q_0E)$

$\Delta U = -W_{Efield} = |F_E|d$

$\Delta V = \Delta U/q_0$

F_g on m is down

$W_{Gfield} = -|F_g|h \quad (F_g = mg)$

$\Delta U_G = -W_{Gfield} = |F_g|h$

$\Delta V_G = \Delta U_G/m$



Parallel Plates

- Releasing a positive test charge from rest at point B...



Electric Potential Energy (conservation of energy ideas)



Work is done to move the charge, so we store potential energy, U_E



Charge is released and energy is converted from U_E to KE



Only the displacement in the direction of the E field matters

(ΔU_E independent of path)

Problem: closed loop path, ABCA

- Work done is path independent
  - Only the initial and final position matter
  - Look for an easy solution!



Problem: find V's and ΔV's



Higher V

lower V

$$|\Delta V_{AC}| = Ed = \left(15\frac{V}{m}\right)(0.25\,m) = 3.75\,V$$

$$\Delta U = q\Delta V = \left(-1.6\times10^{-19}C\right)(3.75\,V) = -6\times10^{-19}\,J$$

Problem: find V and ΔV

$V_1 - V_5 = 3.75$ V

$V_1 = 3.75$ V
$V_2 = 2.8125$ V
$V_3 = 1.875$ V
$V_4 = 0.9375$ V
$V_5 = 0$ V



Electric Potential Energy U_E

- Building up arrangements of charge
  - Energy required to "build" = ΔU
- Bring a point charge in from infinity
  - like charges requires energy
    - repulsive forces
  - unlike charges give up energy
    - attractive forces

$$W = Fd = qEd$$

and $$E = \frac{kq}{r^2}$$

...are difficult to use since E is not a constant.

Can use:

$$U_{12} = \Delta U_{12} = q_2 \Delta V_{\infty 1}$$

$$V_\infty = 0$$

$$V_1 = k\frac{q_1}{r_{12}}$$



U_E for more than two charges

- Don't double count
- Bring each one in from "infinity"



- Bringing together like charges requires energy (force them together)

- Bringing together un-like charges gives up energy (fall together naturally)

$$U_{12} = k\frac{q_1 q_2}{r_{12}}$$

$$U_{23} = k\frac{q_2 q_3}{r_{23}}$$

$$U_{13} = k\frac{q_1 q_3}{r_{13}}$$

Equipotential Surfaces

- E field is perpendicular to the equipotential surfaces
- The surface of a conductor is an equipotential surface
  - no E field parallel to the surface in *Electrostatics*
  - gradually "match" the boundaries



Equipotential Surfaces

Equipotentials are perpendicular to the E-field lines.

E field points “down hill”





Contours of a map analogy



Lines of equal altitude are like Lines of equal potential.

Net force on a positive test charge will point "down hill" just like net force on a boulder will point down hill

F and E are perpendicular to the circles

$$\Delta V_{ab} = \frac{kq}{r_b} - \frac{kq}{r_a}$$

Analogy with Gravity and hills

E field points "down hill" perpendicular to the circles



(c)



(d)

Field gets stronger closer to the point charge. Don't have to go as far to have the same change in electric potential

$$E_r = -\left(\frac{dV}{dr}\right)$$

Slightly misleading (circles would not be evenly spaced for V~1/r)

Equipotential Surfaces

- Imaginary or real surfaces of constant voltage
  - The surfaces of a conductor are equipotential surfaces
- E field and equipotential surfaces are perpendicular to each other



If a charge moves from A to B along an equipotential surface, then

$$\Delta V_{AB} = 0$$

$$\Delta U_{AB} = q\Delta V_{AB} = 0$$

PHYS2321

Week5: Electric Potential and Exam I

Day 2 Outline

1) Hwk: Ch. Study for Exam I

Ch. 24 on capacitance is next.

2) General Exam info

3) Electric Potential

ΔV, ΔU_E, W, ΔK and Δv in uniform E

Finding E from V(x,y,z) or V(r)

Calculating potential for continuous charge distribs

Notes:

Exam I on Friday. Prepare review questions for Wed.

PDF version of "Review1 for Exam 1" updated online.

Try Ch. 21-23 "Test Bank Practices" online.

Hwk keys 1-4 online.

Return Hwk 3 (mean = 8.65/10)